

Fractal dimension: A complementary diagnostic indicator of osteoporosis to bone mineral density



Qiang Chen^{a,*}, Nirong Bao^b, Qingqiang Yao^{c,*}, Zhi-Yong Li^{a,d}

^a Biomechanics Laboratory, School of Biological Science & Medical Engineering, Southeast University, 210096 Nanjing, China

^b Department of Orthopaedics, Jinling Hospital, Nanjing University, 210093 Nanjing, China

^c Department of Orthopaedic Surgery, Nanjing First Hospital, 210006 Nanjing, China

^d School of Chemistry, Physics and Mechanical Engineering, Queensland University of Technology (QUT), Brisbane, QLD 4001, Australia

ARTICLE INFO

Keywords:

Osteoporosis
Diagnostic indicator
Fractal dimension
Bone mineral density

ABSTRACT

Morbidity of osteoporosis is increasing as the world population grows and ages, but bone mineral density (BMD) as a common-used diagnostic indicator is not omnipotent in predicting the bone fragility. According to the definition of osteoporosis by World Health Organization (WHO), the present hypothesis proposes an additional fractal-dimension indicator less than 3 to measure the structural change of bones, and further to diagnose osteoporotic patients with complications, to which BMD is insensitive. Literature reports support that the fractal dimension is more sensitive than BMD in specific cases. The hypothesis shows a promise not only in improving the accuracy of screening osteoporotic patients as a complementary indicator to BMD, but also in evaluating mechanical properties of osteoporotic bone and bone-repair effect of bone tissue engineering.

Introduction

Osteoporosis is a disease occurring in older people, and its morbidity and mortality increase with the increasing aged population. In accordance with the WHO definition [1], osteoporosis is characterized by two factors: low bone mass and deteriorated micro-architecture (Fig. 1, [2]), which enhance subsequent fracture risk due to bone fragility. The present diagnosis on the osteoporosis mainly relies on the BMD measurement of bone mass, but the measurement on the deteriorated micro-architecture is absent.

Diagnostic limitation of the BMD in clinics

Clinically, BMD value indicating the bone mineralization is often employed to judge the occurrence of osteoporosis (i.e., T scores < -2.5) and predict the risk of osteoporotic fracture (ROF). However, BMD was not sensitive to the vertebral compression fracture (VCF) resulting from osteoporosis in type 2 diabetic women [3], and it failed to predict the ROF between the younger and elder postmenopausal women [4]. Since BMD cannot effectively predict the ROF for the specific cases, new measurement is needed for the diagnosis of osteoporosis.

Diagnostic necessity of structural change of osteoporotic bone

The micro-architecture of human bone is porous and inter-connecting. Despite of the correlation between the BMD and the micro-architecture, bone densitometry or related medical devices examining the patients' BMD tell a little about the structural change of abnormal bone. Moreover, bone fragility is a result of the failed structural adaptation, not just low bone mass, and the failed structural adaptation results from a broken balance of bone remodeling [5]. It is the broken balance that causes the osteoporosis, which features trabecular thinning and re-patterning (the ratio of rod- and plate-like trabecula, [6]). Thus, in terms of the structural change of abnormal bone, diagnosing osteoporosis from the standpoint of bone structure is necessary.

Fractal dimension describing structure of abnormal bone

Fractal geometry is often used to describe irregular porous media, which are characterized by a non-Euclidean fractal dimension, and the fractal dimension is a non-integer. In the fields of medicine, it has been employed to treat the structure of abnormal bone, such as osteoarthritis [7] and osteoporosis [8]. Then, addressing the diagnostic limitation of the BMD, is the fractal dimension able to measure the structural change of osteoporotic bones and overcome the limitation? If so, clinicians can diagnose the osteoporosis and further predict bone fracture risk by

* Corresponding authors.

E-mail addresses: chenq999@gmail.com, cq@seu.edu.cn (Q. Chen), Yaoqingqiang@gmail.com (Q. Yao).




Fig. 1. Morphologies of health and osteoporotic bone [2].




Fig. 2. Complementary diagnosis of fractal dimension and BMD.

combining the proposed fractal indicator with the conventional BMD index, see Fig. 2.

The hypothesis

Considering the complex structure of bones, we hypothesize that the fractal dimension of bone matrix (less than 3) could be a complementary indicator to BMD to diagnose the osteoporosis. This is because the fractal dimension represents the changes of bone microstructure, and the BMD indicates the changes of bone mineralization, thus, with both indicators, screening osteoporosis could be more accurate, see Fig. 2. For osteoporosis, a greater fractal dimension of bone matrix indicates smaller loss of bone (or smaller porosity), while a smaller fractal dimension of bone matrix shows a greater loss of bone (or larger porosity). This can be understood by an extreme case, namely, when porosity in bone tends to 0, the fractal dimension approaches Euclidian dimension of 3.

Evaluation of the hypothesis

Fractal growth was considered as a design principle in biological organisms [9]. As a fundamental parameter of fractal geometry, fractal dimension is used to characterize the bone structure, and some studies evidenced that the fractal dimension could be used as a novel strategy to quantify the structural change of bone [10–13].

Calculation of the fractal dimension of bone

The fractal dimension is usually determined by a box-counting approach [13], and the classical expression is $N \sim \delta^{-D}$, where N is the number of boxes with characteristic size δ used to cover porous media, and D is the fractal dimension. Apparently, D is dependent of the characteristic size δ . However, when calculating bone's D on the basis of medical image, we must bear in mind that an appropriate box size should be selected [7], even though low resolution cannot produce a significantly different fractal dimension of an image (or resolution-independent) to an extent [13]. For example, the appropriate size ranges from 65 μm to 1000 μm for trabecular bone in the proximal femur [11], whereas for subchondral trabecular bone in the severe osteoarthritis of the hip, the size is between 30 μm and 4400 μm . Outside the scale, the

fractal dimension may be not correctly calculated [14]. This may be because the box-counted trabecular bone is fractal within the range [7,10]; otherwise, it is non-fractal.

With the box-counting method applied on few selected 2D radiographical images, the values of the fractal dimension were calculated between 1.05 and 1.84 [7,13]. Some literature performed sample analysis in 3D space and reported that the fractal dimension was between 2.08 and 2.75 [15–18]. However, the examination of osteoporotic bones is always regional, and fractal analysis on few selected 2D radiographical images cannot fully describe the structural change of the examined region, thus, 3D sample analysis is more suitable. Moreover, the increased porosity is negatively correlated to the fractal dimension of solid phase [19]. In this sense, the fractal dimension of the cancellous bone influenced by severe osteoarthritis is smaller than that of the normal [7], and this supports the hypothesis, namely, a smaller fractal dimension of bone matrix shows a greater loss of bone.

Higher sensitivity of the fractal dimension than BMD for postmenopausal women

Compared to BMD, fractal dimension has been used to assess the osteoporosis [20–23]. For example, it was reported that the fractal dimension was superior to the BMD by studying the 2D X-ray images of control and osteoporotic groups [21,22]. In detail, a group of postmenopausal women with osteoporotic vertebral fractures and an age-matched control group of women were discriminated by evaluating the fractal ability, and the result showed that the discrimination was stronger than the commonly-used BMD [21]. This solves the problem of the similar BMD between the younger and older groups of postmenopausal women [4]. Also, the trabecular bone in knee osteoarthritis was analyzed, and it was found that the fractal dimension quantifying the structural changes of the trabecular bone is more sensitive to BMD [22]. Moreover, the statistical significance of the fractal dimension for the mandibular and alveolar bone of the postmenopausal women was found [12,23]. In particular, the relationship between the fractal-related Hurst parameter H and BMD of the vertebrae, hip and wrist, were explicitly discussed, and their independent and complementary information was presented, which indicated that the fractal indicator along with BMD opens a new way to diagnose the osteoporosis [20]. In addition, the accuracy of the fractal analysis on screening the osteoporotic patients could reach 95% [24], and this is higher than the BMD result reported in [25], which identified the undiagnosed low BMD in 200,160 postmenopausal women over fifties. Anyhow, all the above-mentioned evidences support the superiority of the fractal dimension to BMD to diagnose the osteoporosis for postmenopausal women.

Consequences of the hypothesis and discussion

With the evidence by evaluating the applicability of the fractal dimension to distinguish the osteoporotic patients, the hypothesis may be significant to clinicians, due to its non-invasiveness and simplification on the complex bone structure. As a complementary indicator to BMD, we must independently conduct correlation analyses of the fractal dimension and the BMD to osteoporosis. Then, the osteoporosis can be discriminated through combining their correlations. However, under the hypothesis, we cannot identify the cases in which the fractal dimension is more sensitive than the BMD, e.g. osteoporotic patients in type 2 diabetic and postmenopausal women [3,4], and other cases in which the fractal dimension is less sensitive. Moreover, the threshold values of the fractal dimension like T scores of the BMD should be determined. The limitations could be overcome by a large number of cases studies.

Not only for the clinical diagnosis of osteoporosis, the hypothesis could also be used in evaluating the effect of scaffold-guided bone repair or regeneration in tissue engineering. This is achieved by the fractal-dimension compatibility between the after-repaired sites and

their surrounding host bone tissues [26]. Besides, it can also be used to correlate the fractal dimension to the mechanical properties of the osteoporotic bone, such as bone's Young's modulus [18,27] and strength [27], which can be non-invasively obtained by existing imaging technologies. These materials parameters are very important for the global assessment for bone-related diseases, in particular for the osteoporosis.

Competing interests

The authors declare that they have no conflict of interest.

Funding

This work was partially supported by the Natural Science Foundation of China (NSFC) (Nos. 31300780, 11422222, 11772093), and the Fundamental Research Funds for the Central Universities (No. 2242016R30014). YQ would like to thank the Key Research Program of Science & Technology Support Program of Jiangsu Province (BE2015613, BE2016763)

References

- [1] Assessment of fracture risk and its application to screening for postmenopausal osteoporosis: report of a WHO study group. WHO Tech Rep Ser 1994;843:1–129.
- [2] <https://www.badbacks.com.au/info/health-information/osteoporosis>.
- [3] Yamamoto M, Yamaguchi T, Yamauchi M, Kaji H, Sugimoto T. Bone mineral density is not sensitive enough to assess the risk of vertebral fractures in type 2 diabetic women. *Calcified Tissue Int* 2007;80(6):353–8.
- [4] Siris ES, Brennan SK, Miller PD, et al. Predictive value of low BMD for 1-Year fracture outcomes is similar for postmenopausal women ages 50–64 and 65 and older: results from the national osteoporosis risk assessment (NORA). *J Bone Miner Res* 2004;19(8):1215–20.
- [5] Seeman E, Delmas PD. Bone quality—the material and structural basis of bone strength and fragility. *N Engl J Med* 2006;354(21):2250–61.
- [6] Wang J, Stein EM, Zhou B, et al. Deterioration of trabecular plate-rod and cortical micro-architecture and reduced bone stiffness at distal radius and tibia in postmenopausal women with vertebral fractures. *Bone* 2016;88:39–46.
- [7] Fazzalari NL, Parkinson IH. Fractal properties of subchondral cancellous bone in severe osteoarthritis of the hip. *J Bone Miner Res* 1997;12(4):632–40.
- [8] Southard TE, Southard KA, Jakobsen JR, Hillis SL, Najim CA. Fractal dimension in radiographic analysis of alveolar process bone. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1996;82(5):569–76.
- [9] Weibel ER. Fractal geometry: a design principle for living organisms. *Am J Physiol* 1991;261(6):L361–9.
- [10] Chung HW, Chu CC, Underweiser M, Wehrli FW. On the fractal nature of trabecular structure. *Med Phys* 1994;21(10):1535–40.
- [11] Fazzalari NL, Parkinson IH. Fractal dimension and architecture of trabecular bone. *J Pathol* 1996;178(1):100–5.
- [12] Oliveira ML, Pedrosa EFNC, Cruz AD, Haiter-Neto F, Paula FJA, Watanabe PCA. Relationship between bone mineral density and trabecular bone pattern in postmenopausal osteoporotic Brazilian women. *Clin Oral Invest* 2013;17(8):1847–53.
- [13] Osman D, Newitt D, Gies A, et al. Fractal based image analysis of human trabecular bone using the box counting algorithm: impact of resolution and relationship to standard measures of trabecular bone structure. *Fractals* 1998;6(03):275–83.
- [14] Kaye BH. The dangerous art of extrapolation for predicting physical phenomena. In: Encarnação JL, editor. *A random walk through fractal dimensions*. Weinheim, Germany: VCH Publishers; 1989. p. 20.
- [15] Wlczek P, Odgaard A, Sernetz M. Fractal 3D analysis of blood vessels and bones. In: Encarnação JL, editor. *Fractal geometry and computer graphics*. Berlin: Springer; 1992. p. 240.
- [16] Wilding RJC, Slabbert JCG, Kathree H, Owen CP, Crombie K, Delpont P. The use of fractal analysis to reveal remodeling in human alveolar bone following the placement of dental implants. *Arch Oral Biol* 1995;40(1):61–72.
- [17] Joo YI, Sone T, Fukunaga M, Lim SG, Onodera S. Effects of endurance exercise on three-dimensional trabecular bone micro-architecture in young growing rats. *Bone* 2003;33(4):485–93.
- [18] Sanchez-Molina D, Velazquez-Ameijide J, Quintana V, et al. Fractal dimension and mechanical properties of human cortical bone. *Med Eng Phys* 2013;35(5):576–82.
- [19] Dathe A, Thullner M. The relationship between fractal properties of solid matrix and pore space in porous media. *Geoderma* 2005;129(3–4):279–90.
- [20] Benhamou CL, Harba R, Lespessailles E, Jacquet G, Tourliere D, Jennane R. Changes in fractal dimension of trabecular bone in osteoporosis: a preliminary study. *Fractals in Biology and Medicine*. Basel: Birkhäuser; 1994. p. 292–9.
- [21] Pothuaud L, Lespessailles E, Harba R, et al. Fractal analysis of trabecular bone texture on radiographs: discriminant value in postmenopausal osteoporosis. *Osteoporos Int* 1998;8(6):618–26.
- [22] Messent EA, Buckland-Wright JC, Blake GM. Fractal analysis of trabecular bone in knee osteoarthritis (OA) is a more sensitive marker of disease status than bone mineral density (BMD). *Calcified Tissue Int* 2005;76(6):419–25.
- [23] Kavitha M, Khan M, Vijayalakshmi KR. Fractal dimension analysis in digital periapical radiographs: a diagnostic indicator of osteoporosis in post-menopausal women. *J Indian Acad Oral Med Radiol* 2017;29(2):84.
- [24] Tafraouti A, El Hassouni M, Toumi H, Lespessailles E, Jennane R. Osteoporosis diagnosis using fractal analysis and support vector machine. *Signal-Image Technology and Internet-Based Systems (SITIS)*, 2014 tenth international conference on. IEEE. 2014. p. 73–7.
- [25] Siris ES, Miller PD, Barrett-Connor E, et al. Identification and fracture outcomes of undiagnosed low bone mineral density in postmenopausal women: results from the National Osteoporosis Risk Assessment. *JAMA* 2001;286(22):2815–22.
- [26] Oshida Y, Hashem A, Nishihara T, Yapchulay MV. Fractal dimension analysis of mandibular bones: toward a morphological compatibility of implants. *Biomed Mater Eng* 1994;4(5):397–407.
- [27] Majumdar S, Lin J, Link T, et al. Fractal analysis of radiographs: assessment of trabecular bone structure and prediction of elastic modulus and strength. *Med Phys* 1999;26(7):1330–40.